arXiv:math/0611211v1 [math.CA] 8 Nov 2006

ON THE CHAOTIC ASYMPTOTICS OF *RAMANUJAN*'S ENTIRE FUNCTION $A_q(z)$

RUIMING ZHANG

Abstract. We will use a discrete analogue of the classical *Laplace* method to show that for infinitely many positive integers n, the main term of the asymptotic expansions of scaled confluent basic hypergeometric functions, including the Ramanujan's entire function $A_q(z)$, could be expressed in θ-functions, and the error term depends on the ergodic property of certain real scaling parameter.

1. Introduction

In [17] we studied some *Plancherel-Rotach* type asymptotics for three families of q-orthogonal polynomials with logarithmic scalings. The asymptotics reveals a remarkable pattern which is quite a contrast to the classical *Plancherel-Rotach* type asymptotics[6, 24]. For each family of such polynomials, we found that when a scaling parameter is above certain critical value, the main term of the asymptotics involves *Ramanujan*'s entire function $A_q(z)$, and the error term is exponentially small; when the scaling parameter is below the critical value, if it is rational, then the main term of the asymptotics has a factor of theta function and the error term is exponentially small, if the scaling parameter is irrational, then the main term also contains some theta function, but we could only show that the error term is no worse than $\mathcal{O}(n^{-1}\log n)$ there.

In this paper, we will use the same technique to study a *Plancherel-Rotach* type asymptotics for the confluent basic hypergeometric series, and we will demonstrate that the asymptotics follows a similar pattern to the case when a scaling parameter is less than some critical value[17], with the only exception that now the base of the theta function explicitly depends on the type of the confluent basic hypergeometric series. Furthermore, the error term, in the irrational scaling parameter case, is closely related to the ergodic property of the number. Generally speaking, we could improve the error term to $\mathcal{O}(n^{-r})$, where r is any the positive number less than the chaotic index(we will define it later in section 2) of the number. In particular, when the scaling parameter is a real algebraic number of degree l, the error term could not be better than $\mathcal{O}(n^{-r})$, where the number r is bounded above by $l-1$, and when the scaling parameter is a *Liouville* number, the error term is better than $\mathcal{O}(n^{-r})$ for any positive number r.

Date: November 6, 2006.

1991 *Mathematics Subject Classification.* Primary 33D45. Secondary 33E05.

Key words and phrases. Plancherel-Rotach asymptotics, θ-function, *q-Airy* function, *Ramanujan*'s entire function, confluent basic hypergeometric series, discrete *Laplace* method, *Chebyshev*'s theorem, *Liouville* number, *Liouville*'s theorem, *Liouville-Roth* constant, irrationality measure, chaotic index of an irrational number.

In section 2, we first introduce the *Ramanujan*'s entire function $A_q(z)$, then define the chaotic index of an irrational number which is related to its ergodic property. In section 3, we will prove asymptotics formulas for $A_q(z)$ in detail, and then just state the results for the confluent basic hypergeometric series without proofs because the proofs are similar to the corresponding cases for $A_q(z)$.

2. Preliminaries

2.1. ***Ramanujan*'s Entire Function.**

Given an arbitrary positive real number $0 < q < 1$ and an arbitrary complex number a, we define[8, 12]

$$(a;q)_\infty = \prod_{k=0}^{\infty}(1-aq^k) \tag{1}$$

and the q-shifted factorials

$$(a;q)_n = \frac{(a;q)_\infty}{(aq^n;q)_\infty}, \tag{2}$$

for any integer $n \in \mathbb{Z}$. Assuming that $|z| < 1$, the q-binomial theorem is[8, 12]

$$\frac{(az;q)_\infty}{(z;q)_\infty} = \sum_{k=0}^{\infty}\frac{(a;q)_k}{(q;q)_k}z^k. \tag{3}$$

Its limiting case

$$(z;q)_\infty = \sum_{k=0}^{\infty}\frac{q^{k(k-1)/2}}{(q;q)_k}(-z)^k, \tag{4}$$

is one of many q-exponentials. For any $n \in \mathbb{N}$, let

$$R_1(a;n) := (aq^n;q)_\infty - 1 \tag{5}$$

and

$$R_2(a;n) := \frac{1}{(aq^n;q)_\infty} - 1, \tag{6}$$

then it is not hard to see that[17]

$$|R_1(a;n)| \le \frac{(-aq^2;q)_\infty aq^n}{1-q} \tag{7}$$

for $a > 0$ and

$$|R_2(a;n)| \le \frac{aq^n}{(1-q)(aq;q)_\infty} \tag{8}$$

for $0 < aq < 1$.

The *Ramanujan*'s entire function $A_q(z)$ is defined as[12]

$$A_q(z) = \sum_{k=0}^{\infty}\frac{q^{k^2}}{(q;q)_k}(-z)^k. \tag{9}$$

Clearly, it satisfies the following three term recurrence

$$A_q(z) - A_q(qz) = qzA_q(q^2z). \tag{10}$$

The function $A_q(z)$ appears repeatedly in Ramanujan's work starting from the Rogers-Ramanujan identities where $A_q(-1)$ and $A_q(-q)$ are expressed as infinite products[2]. It is known that $A_q(z)$ has infinitely many zeros, and all of them are positive[12]. There have been many researches around $A_q(z)$ recently, for example, to properties of and conjectures about its zeros, [4], [5], [9], [15]. The *Ramanujan*'s $A_q(z)$ is also called as *q-Airy* function in the literature, but this is an unfortunate name, since $A_q(z)$ has nothing to do with the classical *Airy* functions.

It is clear that

$$\left|\frac{(1-q)^k}{(q;q)_k} q^{k^2}(-z)^k\right| \le \frac{(q\,|z|)^k}{k!} \tag{11}$$

for $k = 0, 1, \dots$ and for any complex number z. Applying the *Lebesgue*'s dominated convergent theorem we have

$$\lim_{q\to 1} A_q((1-q)z) = e^{-z} \tag{12}$$

for any $z \in \mathbb{C}$. From (11) we also have obtained the inequality

$$|A_q((1-q)z)| \le e^{q|z|} \tag{13}$$

for any complex number z.

For any nonzero complex number z, we define the theta function as

$$\theta(z;q) = \sum_{k=-\infty}^{\infty} q^{k^2/2} z^k, \tag{14}$$

the *Jacobi*'s triple product formula says that[8, 12]

$$\sum_{k=-\infty}^{\infty} q^{k^2/2} z^k = (q, -q^{1/2}z, -q^{1/2}/z; q)_\infty. \tag{15}$$

2.2. **Chaotic Index.** For a positive irrational number θ, *Chebyshev*'s theorem or its more accurate counterpart *Khinchin*'s theorem implies that for any real number β , there exist infinitely many pairs of integers n and m with $n > 0$ such that[10]

$$n\theta = m + \beta + \gamma_n \quad \text{with} \quad |\gamma_n| \le 3/n. \tag{16}$$

Clearly, *Chebyshev*'s theorem and *Khinchin*'s theorem are saying that the arithmetic progression $\{n\theta\}_{n\in\mathbb{Z}}$ is ergodic in $\mathbb{R}$.

Definition 2.1. Given real number θ, the *chaotic index* $\omega(\theta)$ of θ is defined as the least upper bound of the set of real numbers r such that for any real number β, there exist infinitely many integers m and n with $n > 0$ such that

$$|n\theta - \beta - m| \le \frac{C(\theta, r)}{n^r}, \tag{17}$$

where $C(\theta, r)$ is a positive constant which depends only on θ and r.

Theorem 2.2. *The chaotic index $\omega(\theta) \ge 1$ for any irrational number θ .*

Proof. This is a direct consequence of the *Chebyshev*'s theorem or the *Khintchin*'s theorem. □

Recall that the *irrationality measure* (or *Liouville-Roth constant*) $\mu(\theta)$ of a real number θ is defined as the least upper bound of the set of real numbers r such that[28]

$$0 < |n\theta - m| < \frac{1}{n^{r-1}} \tag{18}$$

is satisfied by infinite number of integer pairs (n, m) with $n > 0$. It is clear that we have $\omega(\theta) \leq \mu(\theta) - 1$, since given a real number $r < \omega(\theta)$ in the definition2.1, taking $\beta = 0$ in (17), we have

$$|n\theta - m| \leq \frac{C(\theta, r)}{n^r} \tag{19}$$

Recall that a real algebraic number θ of degree l if it is a root of an irreducible polynomial of degree l over the integer ring. *Liouville*'s theorem in number theory says that for a real algebraic number θ of degree l, there exists a positive constant $K(\theta)$ such that for any integer m and $n > 0$ we have

$$|n\theta - m| > \frac{K(\theta)}{n^{l-1}}. \tag{20}$$

A *Liouville* number is a real number θ such that for any positive integer l there exist infinitely many integers n and m with $n > 1$ such that[28]

$$0 < |n\theta - m| < \frac{1}{n^{l-1}}, \tag{21}$$

and the terms in the continued fraction expansion of every *Liouville* number are unbounded. Even though the set of all *Liouville* numbers is of Lebesgue measure zero, it is known that almost all real numbers are *Liouville* numbers topologically.

Theorem 2.3. *The chaotic index ω has the following properties:*

1. *For any rational number $\theta \in \mathbb{Q}$, its chaotic index $\omega(\theta)$ is zero;*
2. *For any real algebraic number θ of degree l, its chaotic index $\omega(\theta) \leq l - 1$;*
3. *For any Liouville number θ, its chaotic index $\omega(\theta) = \infty$.*

Proof. Let $\theta = \frac{a}{b}$ for some $a, b \in \mathbb{N}$ with $(a, b) = 1$ and $b > 0$. If the first claim is not true, then there must be a positive number σ, and infinitely many pair of integers m and n with $n > 0$ such that

$$|n\theta - m - \beta| \leq \frac{C(\theta)}{n^\sigma}.$$

for all $\beta \in \mathbb{R}$. Since $n\theta - m$ are rational numbers with remainders in the set $\{0, \frac{1}{b}, \ldots, \frac{b-1}{b}\}$, so we could pick a real number β and find some positive constant $M(b, \beta)$ such that

$$|n\theta - m - \beta| \geq M(b, \beta) > 0,$$

for any integer pairs n and m with $n > 0$, which implies that

$$M(b, \beta) \leq \frac{C(\theta)}{n^\sigma}.$$

The last inequality is obviously false for n is large enough, which proves the first assertion. The second and the third assertions follow from the definition 2.1. □

It is clear that the quadratic irrationals such as $\sqrt{2}$ have *chaotic index* $\omega = 1$. A thorough study on the *chaotic index* $\omega(\theta)$ as a function in variable θ is very desirable, but it is out of the scope of the current paper. It is also very natural for us to conjecture the following:

Conjecture. *The chaotic index $\omega(\theta) = \mu(\theta) - 1$ for any real number θ and $\omega(\theta) = l - 1$ if θ is a real algebraic number of degree l.*

3. Main Results

For any positive real number t, we consider the following set

$$\mathbb{S}(t) = \{\{nt\} : n \in \mathbb{N}\}. \tag{22}$$

It is clear that $\mathbb{S}(t) \subset [0,1)$ and it is a finite set when t is rational. In this case, for any $\lambda \in \mathbb{S}(t)$, there are infinitely many pair of integers n and m with $n > 0$ such that

$$nt = m + \lambda, \tag{23}$$

where

$$m = \lfloor nt \rfloor. \tag{24}$$

If t is a positive irrational number, then $\mathbb{S}(t)$ is a subset of $(0,1)$ with infinite elements, and it is well-known that $\mathbb{S}(t)$ is uniformly distributed in $(0,1)$. Let r be any positive number less the chaotic index $\omega(t)$, for any $\beta \in [0,1)$, there are infinitely many pair of integers n and m with $n > 0$ such that

$$nt = m + \beta + \gamma_n \tag{25}$$

with

$$|\gamma_n| \le \frac{C(t,r)}{n^r}, \tag{26}$$

for some constant $C(t,r)$. It is clear that for n large enough, we may also have

$$m = \lfloor nt \rfloor. \tag{27}$$

In the statements of our results, we also need the arithmetic function $\chi(n)$, the principal character modulo 2, which is defined as

$$\chi(n) = \begin{cases} 1 & 2 \nmid n \\ 0 & 2 \mid n \end{cases}. \tag{28}$$

In the proof of the irrational scaling parameter case, we also make use of the elementary inequality

$$|e^x - 1| \le |x| e^{|x|} \tag{29}$$

for all $x \in \mathbb{C}$.

3.1. Asymptotics of $A_q(z)$.

For the *Ramanujan*'s entire function $A_q(z)$, we have the following:

Theorem 3.1. *Given an arbitrary positive real number $0 < q < 1$ and an arbitrary nonzero complex number u, we have*

1. *Assuming that t is a positive rational number, for any $\lambda \in \mathbb{S}(t)$, there are infinitely many pairs of integers n and m satisfying (23) and (24) with $n > 0$. For these n and m we have*

$$A_q(q^{-nt}u) = \frac{(-u)^{\lfloor m/2 \rfloor}\left\{\theta(-u^{-1}q^{\chi(m)+\lambda}; q^2) + r(n)\right\}}{(q;q)_\infty q^{\lfloor m/2 \rfloor (nt - \lfloor m/2 \rfloor)}} \tag{30}$$

with

$$|r(n)| \le \frac{3(-q^3;q)_\infty \theta(|u|^{-1} q^{\chi(m)+\lambda};q^2)}{(q;q)_\infty} \times \left\{ q^{nt/4} + \frac{q^{m^2/16}}{|u|^{\lfloor m/4\rfloor+1}} \right\}. \tag{31}$$

(2) *Assuming that t is a positive irrational number and r is a positive number strictly less the chaotic index $\omega(t)$ of t, for any real number $\beta \in [0,1)$, there are infinitely many pair of integers n and m with $n > 0$ satisfying (25), (26) and (27). For such n and m we have*

$$A_q(q^{-nt}u) = \frac{(-u)^{\lfloor m/2\rfloor} \left\{ \theta(-u^{-1}q^{\chi(m)+\beta};q^2) + \mathcal{O}\left(n^{-r}\right) \right\}}{(q;q)_\infty q^{\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}}. \tag{32}$$

when n is large enough.

Proof. From (9), we have

$$A_q(q^{-nt}u) = \sum_{k=0}^{\infty} \frac{q^{k^2-knt}}{(q;q)_k}(-u)^k. \tag{33}$$

In the case that t is a positive rational number, for any $\lambda \in \mathbb{S}(t)$ and n, m are large, we have

$$\begin{aligned} A_q(q^{-nt}u)(q;q)_\infty &= \sum_{k=0}^{\infty} (q^{k+1};q)_\infty q^{k^2-km-k\lambda}(-u)^k \\ &= s_1 + s_2 \end{aligned} \tag{34}$$

where

$$s_1 = \sum_{k=0}^{\lfloor m/2\rfloor} (q^{k+1};q)_\infty q^{k^2-km-k\lambda}(-u)^k \tag{35}$$

and

$$s_2 = \sum_{k=\lfloor m/2\rfloor+1}^{\infty} (q^{k+1};q)_\infty q^{k^2-km-k\lambda}(-u)^k. \tag{36}$$

In s_1 we change the summation from k to $\lfloor m/2\rfloor - k$, we have

$$\begin{aligned}
\frac{s_1 q^{\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}}{(-u)^{\lfloor m/2\rfloor}} &= \sum_{k=0}^{\lfloor m/2\rfloor} (q^{\lfloor m/2\rfloor-k+1};q)_\infty q^{k^2}(-u^{-1}q^{\chi(m)+\lambda})^k \\
&= \sum_{k=0}^{\infty} q^{k^2}(-u^{-1}q^{\chi(m)+\lambda})^k \\
&- \sum_{k=\lfloor m/4\rfloor+1}^{\infty} q^{k^2}(-u^{-1}q^{\chi(m)+\lambda})^k \\
&+ \sum_{k=0}^{\lfloor m/4\rfloor} q^{k^2}(-u^{-1}q^{\chi(m)+\lambda})^k \left((q^{\lfloor m/2\rfloor-k+1};q)_\infty - 1\right) \\
&+ \sum_{k=\lfloor m/4\rfloor+1}^{\lfloor m/2\rfloor} (q^{\lfloor m/2\rfloor-k+1};q)_\infty q^{k^2}(-u^{-1}q^{\chi(m)+\lambda})^k \\
&= \sum_{k=0}^{\infty} q^{k^2}(-u^{-1}q^{\chi(m)+\lambda})^k + s_{11} + s_{12} + s_{13}.
\end{aligned}$$

Clearly,

$$\begin{aligned}
|s_{11}+s_{13}| &\le 2\sum_{k=\lfloor m/4\rfloor+1}^{\infty} q^{k^2}(|u|^{-1}q^{\chi(m)+\lambda})^k \\
&\le \frac{2q^{m^2/16}}{|u|^{\lfloor m/4\rfloor+1}}\theta(|u|^{-1}q^{\chi(m)+\lambda};q^2).
\end{aligned}$$

By (7), for $0 \le k \le \lfloor m/4\rfloor$, we have

$$\left|(q^{\lfloor m/2\rfloor-k+1};q)_\infty - 1\right| = \frac{(-q^3;q)_\infty}{(q;q)_\infty}q^{nt/4}, \tag{37}$$

then

$$\begin{aligned}
|s_{12}| &\le \frac{(-q^3;q)_\infty q^{nt/4}}{(q;q)_\infty}\sum_{k=0}^{\infty} q^{k^2}(|u|^{-1}q^{\chi(m)+\lambda})^k \\
&\le \frac{q^{nt/4}(-q^3;q)_\infty\theta(|u|^{-1}q^{\chi(m)+\lambda};q^2)}{(q;q)_\infty},
\end{aligned}$$

hence

$$\frac{s_1 q^{\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}}{(-u)^{\lfloor m/2\rfloor}} = \sum_{k=0}^{\infty} q^{k^2}(-u^{-1}q^{\chi(m)+\lambda})^k + r_1(n) \tag{38}$$

with

$$\begin{aligned}
|r_1(n)| &\le \frac{2(-q^3;q)_\infty\theta(|u|^{-1}q^{\chi(m)+\lambda};q^2)}{1-q} \\
&\times \left\{q^{nt/4} + \frac{q^{m^2/16}}{|u|^{\lfloor m/4\rfloor+1}}\right\}.
\end{aligned} \tag{39}$$

In s_2 we change the summation from k to $k+\lfloor m/2\rfloor$

$$\begin{aligned}\frac{s_2 q^{\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}}{(-u)^{\lfloor m/2\rfloor}} &= \sum_{k=1}^{\infty}(q^{\lfloor m/2\rfloor+k+1};q)_\infty q^{k^2}(-uq^{-\chi(m)-\lambda})^k \\ &= \sum_{k=1}^{\infty} q^{k^2}(-uq^{-\chi(m)-\lambda})^k \\ &+ \sum_{k=1}^{\infty} q^{k^2}(-uq^{-\chi(m)-\lambda})^k\left[(q^{\lfloor m/2\rfloor+1};q)_\infty - 1\right] \\ &= \sum_{k=-\infty}^{-1} q^{k^2}(-u^{-1}q^{\chi(m)+\lambda})^k + r_2(n).\end{aligned}$$

By (7), for $k\ge 1$

$$\left|(q^{\lfloor m/2\rfloor+1};q)_\infty - 1\right| \le \frac{q^{nt/2}(-q^3;q)_\infty}{(q;q)_\infty}, \tag{40}$$

then

$$|r_2(n)| \le \frac{q^{nt/2}(-q^3;q)_\infty\theta(|u|^{-1}q^{\chi(m)+\lambda};q^2)}{(q;q)_\infty}. \tag{41}$$

Thus we have proved that

$$A_q(q^{-nt}u) = \frac{(-u)^{\lfloor m/2\rfloor}\left\{\theta(-u^{-1}q^{\chi(m)+\lambda};q^2)+r(n)\right\}}{(q;q)_\infty q^{\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}} \tag{42}$$

with

$$\begin{aligned}|r(n)| &\le \frac{3(-q^3;q)_\infty\theta(|u|^{-1}q^{\chi(m)+\lambda};q)}{1-q} \\ &\times \left\{q^{nt/4} + \frac{q^{m^2/16}}{|u|^{\lfloor m/4\rfloor+1}}\right\}.\end{aligned} \tag{43}$$

for n, m and λ satisfying (23) , (24)with n and m are large enough.

In the case that t is a positive irrational number, for any real number $\beta\in[0,1)$, when n and m satisfy (25), (26) and (27) and n is large enough, we take

$$\nu_n = \left\lfloor -\frac{\log n}{\log q}\right\rfloor, \tag{44}$$

then we have

$$\begin{aligned}A_q(q^{-nt}u)(q;q)_\infty &= \sum_{k=0}^{\infty}(q^{k+1};q)_\infty q^{k^2-km-k\beta-k\gamma_n}(-u)^k \\ &= s_1+s_2,\end{aligned} \tag{45}$$

with

$$s_1 = \sum_{k=0}^{\lfloor m/2\rfloor}(q^{k+1};q)_\infty q^{k^2-km-k\beta-k\gamma_n}(-u)^k \tag{46}$$

and

$$s_2 = \sum_{k=\lfloor m/2\rfloor+1}^{\infty} (q^{k+1};q)_\infty q^{k^2-km-k\beta-k\gamma_n}(-u)^k. \tag{47}$$

In s_1, we change summation from k to $\lfloor m/2\rfloor - k$,

$$\begin{aligned}
\frac{s_1 q^{\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}}{(-u)^{\lfloor m/2\rfloor}} &= \sum_{k=0}^{\lfloor m/2\rfloor} (q^{\lfloor m/2\rfloor-k+1};q)_\infty q^{k^2}(-u^{-1}q^{\chi(m)+\beta+\gamma_n})^k \\
&= \sum_{k=0}^{\infty} q^{k^2}(-u^{-1}q^{\chi(m)+\beta})^k \\
&- \sum_{k=\nu_n+1}^{\infty} q^{k^2}(-u^{-1}q^{\chi(m)+\beta})^k \\
&+ \sum_{k=0}^{\nu_n} q^{k^2}(-u^{-1}q^{\chi(m)+\beta})^k \left(q^{k\gamma_n}-1\right) \\
&+ \sum_{k=0}^{\nu_n} q^{k^2}(-u^{-1}q^{\chi(m)+\beta+\gamma_n})^k \left\{(q^{\lfloor m/2\rfloor-k+1};q)_\infty - 1\right\} \\
&+ \sum_{k=\nu_n+1}^{\lfloor m/2\rfloor} q^{k^2}(-u^{-1}q^{\chi(m)+\beta+\gamma_n})^k (q^{\lfloor m/2\rfloor-k+1};q)_\infty \\
&= \sum_{k=0}^{\infty} q^{k^2}(-u^{-1}q^{\chi(m)+\beta})^k + s_{11}+s_{12}+s_{13}+s_{14},
\end{aligned}$$

thus there exists some constant $c_{11}(q,u)$ such that

$$\begin{aligned}
|s_{11}+s_{14}| &\le 2\sum_{k=\nu_n+1}^{\infty} q^{k^2}(|u|^{-1}q^{\chi(m)+\beta})^k \\
&\le c_{11}(q,u)n^{-r}\log n
\end{aligned}$$

for n large enough, and there exists some constant $c_{12}(q,u)$ such that

$$|s_{12}+s_{13}| \le c_{12}(q,u)n^{-r}\log n,$$

for n large enough. Thus for n, m and β satisfying (25) , (26) and (27) and n is large enough, there exists some constant $c_1(q,u)$ such that

$$\frac{s_1 q^{\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}}{(-u)^{\lfloor m/2\rfloor}} = \sum_{k=0}^{\infty} q^{k^2}(-u^{-1}q^{\chi(m)+\beta})^k + e_1(n) \tag{48}$$

with

$$|e_1(n)| \le c_1(q,u)n^{-r}\log n. \tag{49}$$

Similarly, in s_2 we change summation from k to $k+\lfloor m/2\rfloor$, and we could show that there for n, m and β satisfying (25) , (26) and (27) and n is large enough, there exists some constant $c_2(q,u)$ such that

$$\frac{s_2 q^{\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}}{(-u)^{\lfloor m/2\rfloor}} = \sum_{k=-\infty}^{-1} q^{k^2}(-u^{-1}q^{\chi(m)+\beta})^k + e_2(n) \tag{50}$$

with

$$|e_2(n)| \le c_2(q,u)n^{-r}\log n. \tag{51}$$

Thus we have

$$A_q(q^{-nt}u) = \frac{(-u)^{\lfloor m/2\rfloor}\left\{\theta(-u^{-1}q^{\chi(m)+\beta};q^2) + \mathcal{O}\left(n^{-r}\log n\right)\right\}}{(q;q)_\infty q^{\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}} \tag{52}$$

for n, m and β satisfying (25) , (26) and (27) and n is large enough. Since $\log n = \mathcal{O}(n^{-\epsilon})$ for any positive number ϵ, and any r is an arbitrary positive number strictly less the chaotic index $\omega(t)$, thus we have proved the last assertion. □

3.2. **Asymptotics of Confluent Basic Hypergeometric Functions.** The phenomenon demonstrated with $A_q(z)$ in theorem3.1 is universal for a class of entire basic hypergeometric function of type

$$f(z) = \sum_{k=0}^{\infty}\frac{(a_1,\dots,a_r;q)_k q^{lk^2}}{(b_1,\dots,b_s;q)_k}z^k, \tag{53}$$

with $l > 0$, where

$$(a_1,\dots,a_r;q)_k = \prod_{j=1}^{r}(a_j;q)_k. \tag{54}$$

A basic hypergeometric series is formally defined as

$${}_r\phi_s\left(\begin{array}{c} a_1,\dots,a_r \\ b_1,\dots,b_s\end{array}\middle| q,z\right) = \sum_{k=0}^{\infty}\frac{(a_1,\dots,a_r;q)_k}{(b_1,\dots,b_s;q)_k}z^k\left(-q^{(k-1)/2}\right)^{k(s+1-r)}, \tag{55}$$

when $s+1-r>0$, it is clear that the function

$${}_r\phi_s\left(\begin{array}{c} a_1,\dots,a_r \\ b_1,\dots,b_s\end{array}\middle| q,-zq^{(s+1-r)/2}\right) \tag{56}$$

is of the form (53) with

$$l = \frac{s+1-r}{2}. \tag{57}$$

For a clean statement of the following theorem, we also define

$$c(r,s;q) := \frac{(a_1,\dots,a_r;q)_\infty}{(b_1,\dots,b_s;q)_\infty}. \tag{58}$$

We have similar results for (53) as theorem 3.1 for $A_q(z)$, and their proof is very similar to the proof of theorem 3.1 without any significant changes.

Theorem 3.2. *Assume that*

$$0 \le a_1,\dots,a_r,b_1,\dots,bs < 1, \tag{59}$$

and $0<q<1$*. For any nonzero complex number* u*, we have*

(1) *Given any positive rational number* t*, let* $\lambda \in \mathbb{S}(t)$*, there are infinitely many pair of integers* n *and* m *satisfying (23) and (24) with* $n>0$*. For these* n *and* m *we have*

$$f(q^{-lnt}u) = \frac{(-u)^{\lfloor m/2\rfloor}\left\{\theta(u^{-1}q^{l\chi(m)+l\lambda};q^{2l}) + r(n)\right\}}{c(s,r;q)q^{l\lfloor m/2\rfloor(nt-\lfloor m/2\rfloor)}} \tag{60}$$

with

$$
\begin{aligned}
|r(n)| \quad &\leq \quad \frac{2^{r+s+2}\theta(|u|^{-1}\, q^{l\chi(m)+l\lambda}; q^{2l})}{(q;q)_\infty^{r+s}} \\
&\times \left\{ \frac{\prod_{j=1}^{s}(-b_j q^2;q)_\infty}{\prod_{j=1}^{r}(a_j q;q)_\infty} q^{nt/4} + \frac{q^{lm^2/16}}{|u|^{\lfloor m/4 \rfloor+1}} \right\}.
\end{aligned}
\tag{61}
$$

for n, m *and* λ *satisfying (23) and (24) with* n *is large enough.*

(2) *Given a positive irrational number* t *and a positive number* $r < \omega(t)$, *for any real number* $\beta \in [0, 1)$, *there are infinitely many pair of integers* n *and* m *satisfying (25), (26) and (27) with* $n > 0$. *For such* n *and* m *we have*

$$
f(q^{-lnt}u) = \frac{(-u)^{\lfloor m/2 \rfloor}\left\{\theta(u^{-1}q^{l\chi(m)+l\lambda}; q^{2l}) + \mathcal{O}\left(n^{-r}\right)\right\}}{c(s,r;q)q^{l\lfloor m/2 \rfloor(nt-\lfloor m/2 \rfloor)}}.
\tag{62}
$$

for n *is large enough.*

REFERENCES

[1] N. I. Akhiezer, The Classical Moment Problem and Some Related Questions in Analysis, English translation, Oliver and Boyed, Edinburgh, 1965.

[2] G. E. Andrews, q-series: Their development and application in analysis, number theory, combinatorics, physics, and computer algebra, CBMS Regional Conference Series, number 66, American Mathematical Society, Providence, R.I. 1986.

[3] G. E. Andrews, Ramanujan's "Lost" Note book VIII: The entire Rogers-Ramnujan function, Advances in Math. 191 (2005), 393–407.

[4] G. E. Andrews, Ramanujan's "Lost" Note book IX: The entire Rogers-Ramnujan function, Advances in Math. 191 (2005), 408–422.

[5] G. E. Andrews, R. A. Askey, and R. Roy, Special Functions, Cambridge University Press, Cambridge, 1999.

[6] P. Deift, Orthogonal Polynomials and Random Matrices: a Riemann-Hilbert Approach, American Mathematical Society, Providence, 2000.

[7] P. Deift, T. Kriecherbauer, K. T-R. McLaughlin, S. Venakides, and X. Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, Comm. Pure Appl. Math. 52 (1999), 1491–1552.

[8] G. Gasper and M. Rahman, Basic Hypergeometric Series, second edition Cambridge University Press, Cambridge, 2004.

[9] W. K. Hayman, On the zeros of a q-Bessel function, Contemporary Mathematics, volume 382, American Mathematical Society, Providence, 2005, 205–216.

[10] Hua Loo Keng, Introduction to Number Theory, Springer-Verlag, Berlin Heidelberg New York, 1982.

[11] M. E. H. Ismail, Asymptotics of q-orthogonal polynomials and a q-Airy function, Internat. Math. Res. Notices 2005 No 18 (2005), 1063–1088.

[12] M. E. H. Ismail, Classical and Quantum Orthogonal Polynomials in one Variable, Cambridge University Press, Cambridge, 2005.

[13] M. E. H. Ismail and X. Li, Bounds for extreme zeros of orthogonal polynomials, Proc. Amer. Math. Soc. 115 (1992), 131–140.

[14] M. E. H. Ismail and D. R. Masson, q-Hermite polynomials, biorthogonal rational functions, Trans. Amer. Math. Soc. 346 (1994), 63–116.

[15] M. E. H. Ismail and C. Zhang, Zeros of entire functions and a problem of Ramanujan, Advances in Math., (2007), to appear.

[16] M. E. H. Ismail and R. Zhang, Scaled asymptotics for q-polynomials, Comptes Rendus, submitted.

[17] M. E. H. Ismail and R. Zhang, Chaotic and Periodic Asymptotics for q-Orthogonal Polynomials, joint with Mourad E.H. Ismail, International Mathematics Research Notices, accepted.

[18] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta, Y. Yamada, Hypergeometric solutions to the q-Painlev\'{e} equations, Internat. Math. Res. Notices 47 (2004), 2497–2521.

[19] R. Koekoek and R. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogues, Reports of the Faculty of Technical Mathematics and Informatics no. 98-17, Delft University of Technology, Delft, 1998.

[20] M. L. Mehta, Random Matrices, third edition, Elsevier, Amsterdam, 2004.

[21] W.-Y. Qiu and R.Wong, Uniform asymptotic formula for orthogonal polynomials with exponential weight, SIAM J. Math. Anal.31 (2000), 992–1029.

[22] S. Ramanujan, The Lost Notebook and Other Unpublished Papers (Introduction by G. E. Andrews), Narosa, New Delhi, 1988.

[23] E. B. Saff and V. Totik, Logarithmic Potentials With External Fields, Springer-Verlag, New York, 1997.

[24] G. Szeg\H{o}, Orthogonal Polynomials, Fourth Edition, Amer. Math. Soc., Providence, 1975.

[25] Z. Wang and R. Wong, Uniform asymptotics for the Stieltjes-Wigert polynomials: the Riemann-Hilbert approach, to appear.

[26] R. Wong, Asymptotic Approximations of Intehrals, Academic Press, Boston, 1989.

[27] E. T. Whittaker and G. N. Watson, A Course of Modern Analysis, fourth edition, Cambridge University Press, Cambridge, 1927. Hua Loo Keng, Introduction to Number Theory, Springer-Verlag, Berlin Heidelberg New York, 1982.

[28] Wikipedia.org, *http://en.wikipedia.org/wiki/Liouville_ number.*

E-mail address: `ruimingzhang@yahoo.com`